# ภาคผนวก

# ภาคผนวก ก

## ผู้เชี่ยวชาญตรวจสอบคุณภาพเครื่องมือ

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

# ภาคผนวก ข

## รายนามผู้เชี่ยวชาญตรวจสอบเครื่องมือในการวิจัย

## รายนามผู้เชี่ยวชาญตรวจสอบเครื่องมือในการวิจัย

รายนามผู้เชี่ยวชาญตรวจสอบเครื่องมือในการวิจัย เพื่อพิจารณาความครอบคลุมของเนื้อหา ความถูกต้องของภาษา พร้อมทั้งข้อเสนอแนะในการปรับปรุงเครื่องมือในการวิจัย มีดังนี้

1. ผู้ช่วยศาตราจารย์ ดร.ฆนัท ธาตุทอง ตำแหน่ง คณบดีบัณฑิตวิทยาลัย มหาวิทยาลัยราชภัฏสงขลา

2. นายแพทย์ธิติพันธ์ ธานีรัตน์ ตำแหน่ง นายแพทย์ชำนาญการ (ผู้เชี่ยวชาญด้านสุขภาพจิตและการพยาบาลจิตเวชผู้ป่วยโรคซึมเศร้า) โรงพยาบาลสวนสราญรมย์ ตำบลท่าข้าม อำเภอพุนพิน จังหวัดสุราษฎร์ธานี

3. แพทย์หญิงจุฬาพันธุ์ เหมกุล ตำแหน่ง นายแพทย์ชำนาญการ โรงพยาบาลจิตเวชสงขลาราชนครินทร์ อำเภอเมือง จังหวัดสงขลา

4. นางอรนลิน สิงขรณ์ ตำแหน่ง พยาบาลวิชาชีพชำนาญการ (พยาบาลปฏิบัติการขั้นสูง (APN) ผู้เชี่ยวชาญด้านสุขภาพจิตและการพยาบาลจิตเวชผู้ป่วยโรคซึมเศร้า) โรงพยาบาลสวนสราญรมย์ ตำบลท่าข้าม อำเภอพุนพิน จังหวัดสุราษฎร์ธานี

5. นายติณพัฒน์ แก้วยอดทิวัตถ์ ตำแหน่ง นักจิตวิทยาคลินิกชำนาญการ โรงพยาบาลจิตเวชสงขลาราชนครินทร์ อำเภอเมือง จังหวัดสงขลา

# ภาคผนวก ค

## หนังสือขอความอนุเคราะห์การพิทักษ์สิทธิ์ของกลุ่มตัวอย่าง

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

# ภาคผนวก ง

## ค่าดัชนีความสอดคล้องของเครื่องมือ

## ค่าดัชนีความสอดคล้องระหว่างข้อคำถามกับลักษณะเฉพาะกลุ่ม จากความคิดเห็นของผู้เชี่ยวชาญ

+1 เมื่อแน่ใจว่าข้อคำถามนั้นเป็นตัวแทนลักษณะเฉพาะกลุ่มนั้น
0 เมื่อไม่แน่ใจว่าข้อคำถามนั้นเป็นตัวแทนลักษณะเฉพาะกลุ่มนั้นหรือไม่
-1 เมื่อแน่ใจว่าข้อคำถามนั้นไม่เป็นตัวแทนลักษณะเฉพาะกลุ่มนั้นคะแนน

| ข้อคำถาม | คะแนนความคิดเห็นของผู้เชี่ยวชาญ (คนที่) | | | | | คะแนนรวม | ค่า IOC | ผลการคัดเลือก |
|---|---|---|---|---|---|---|---|---|
| | 1 | 2 | 3 | 4 | 5 | | | |
| 1 | 1 | -1 | 1 | 1 | 1 | 3 | 0.60 | เลือกไว้ |
| 2 | 1 | 1 | 1 | 1 | 1 | 5 | 1.00 | เลือกไว้ |
| 3 | 1 | 1 | 1 | 1 | 1 | 5 | 1.00 | เลือกไว้ |
| 4 | 1 | -1 | 1 | 1 | 1 | 3 | 0.60 | เลือกไว้ |
| 5 | 1 | 1 | 1 | 0 | 1 | 4 | 0.80 | เลือกไว้ |
| 6 | 1 | 0 | 0 | 1 | 1 | 3 | 0.60 | เลือกไว้ |
| 7 | 1 | 0 | 0 | 1 | 1 | 3 | 0.60 | เลือกไว้ |
| 8 | 1 | 1 | -1 | 1 | 0 | 2 | 0.40 | คัดออก |
| 9 | 1 | -1 | 1 | 0 | 0 | 1 | 0.20 | คัดออก |
| 10 | 1 | 0 | 1 | 1 | 0 | 3 | 0.40 | คัดออก |
| 11 | 1 | 1 | 1 | 1 | 0 | 4 | 0.80 | เลือกไว้ |
| 12 | 1 | 1 | 0 | 1 | -1 | 2 | 0.40 | คัดออก |
| 13 | 1 | -1 | 1 | 1 | 0 | 2 | 0.40 | คัดออก |
| 14 | 1 | 0 | 1 | 1 | 0 | 3 | 0.60 | เลือกไว้ |
| 15 | 1 | 1 | 1 | 1 | -1 | 3 | 0.60 | เลือกไว้ |
| 16 | 1 | -1 | 1 | 1 | -1 | 1 | 0.20 | คัดออก |
| 17 | 1 | 1 | 1 | 0 | -1 | 2 | 0.40 | คัดออก |
| 18 | 1 | -1 | 1 | -1 | -1 | -1 | -0.20 | คัดออก |
| 19 | 1 | 1 | 0 | 1 | -1 | 2 | 0.40 | คัดออก |
| 20 | 1 | 1 | 1 | 1 | 0 | 4 | 0.80 | คัดออก |
| 21 | 1 | 1 | 1 | 1 | 0 | 4 | 0.80 | คัดออก |
| 22 | 1 | -1 | 1 | 1 | 1 | 3 | 0.60 | คัดออก |
| 23 | 1 | 1 | 1 | 0 | 1 | 4 | 0.80 | คัดออก |
| 24 | 1 | -1 | 0 | 1 | 0 | 1 | 0.20 | คัดออก |
| 25 | 1 | 1 | 1 | 1 | 0 | 4 | 0.80 | คัดออก |
| 26 | 1 | 1 | 1 | 1 | 1 | 5 | 1.00 | เลือกไว้ |
| 27 | 1 | -1 | 1 | 0 | -1 | 0 | 0.00 | คัดออก |

**ค่าดัชนีความสอดคล้องระหว่างข้อคำถามกับลักษณะเฉพาะกลุ่ม**
**จากความคิดเห็นของผู้เชี่ยวชาญ (ต่อ)**

| ข้อคำถาม | คะแนนความคิดเห็นของผู้เชี่ยวชาญ (คนที่) | | | | | คะแนนรวม | ค่า IOC | ผลการคัดเลือก |
|---|---|---|---|---|---|---|---|---|
| | 1 | 2 | 3 | 4 | 5 | | | |
| 28 | 1 | -1 | 1 | 0 | -1 | 0 | 0.00 | คัดออก |
| 29 | 1 | -1 | 1 | 1 | 1 | 3 | 0.60 | คัดออก |
| 30 | 1 | 1 | 0 | 0 | 1 | 3 | 0.60 | เลือกไว้ |
| 31 | 1 | 1 | -1 | 1 | -1 | 1 | 0.20 | คัดออก |
| 32 | 1 | 1 | 1 | 0 | -1 | 2 | 0.40 | คัดออก |
| 33 | 1 | -1 | 1 | 1 | -1 | 1 | 0.20 | คัดออก |
| 34 | 1 | 1 | 1 | 1 | 1 | 5 | 1.00 | เลือกไว้ |
| 35 | 1 | 1 | 1 | 1 | -1 | 3 | 0.60 | เลือกไว้ |
| 36 | 1 | 1 | 1 | 1 | -1 | 3 | 0.60 | เลือกไว้ |

SONGKHLA RAJABHAT UNIVERSITY

## ค่าความเที่ยงของเครื่องมือ

**Reliability**

****** Method 1 (space saver) will be used for this analysis ******

–

R E L I A B I L I T Y  A N A L Y S I S  -  S C A L E  (A L P H A)

Reliability Coefficients

N of Cases =  40.0  N of Items = 15

Alpha =  .7210

มหาวิทยาลัยราชภัฏสงขลา SONGKHLA RAJABHAT UNIVERSITY

# ภาคผนวก จ
## เครื่องมือการวิจัย

# เครื่องมือการวิจัย

**เรียน ผู้ให้ข้อมูลการวิจัย**

ดิฉัน นางสุดา ยุทธโท เป็นนักศึกษาปริญญาโทหลักสูตรสาธารณสุขศาสตรมหาบัณฑิต สาขาวิชาสาธารณสุขชุมชน บัณฑิตวิทยาลัย มหาวิทยาลัยราชภัฏสงขลา กำลังศึกษาเกี่ยวกับความความสัมพันธ์ระหว่างความเชื่ออำนาจภายใน-ภายนอกตนด้านสุขภาพกับภาวะซึมเศร้าของผู้ป่วยโรคซึมเศร้า ซึ่งมีวัตถุประสงค์เพื่อศึกษาระดับความเชื่ออำนาจภายใน-ภายนอกตนด้านสุขภาพของผู้ป่วยโรคซึมเศร้าและความสัมพันธ์ระหว่างความเชื่ออำนาจภายใน-ภายนอกตนด้านสุขภาพกับภาวะซึมเศร้าของผู้ป่วยโรคซึมเศร้า นำผลการศึกษาไปใช้ในการส่งเสริมป้องกันปัญหาสุขภาพและให้การดูแลผู้ป่วยโรคซึมเศร้าและประยุกต์ใช้กับผู้ป่วยอื่น ให้เหมาะสมและมีประสิทธิภาพมากยิ่งขึ้น

จึงใคร่ขอความกรุณาท่านในการเข้าร่วมการศึกษาโดยตอบแบบวัดตามความเป็นจริง ซึ่งการเข้าร่วมการศึกษาในครั้งนี้เป็นไปตามความสมัครใจของท่านและหากเมื่อท่านเข้าร่วมการศึกษาแล้วท่านมีสิทธิ์ยกเลิกการเข้าร่วมการศึกษาเมื่อไรก็ได้โดยไม่ต้องบอกเหตุผล ซึ่งจะไม่มีผลกระทบใดๆต่อการรักษาและตัวท่าน ทั้งนี้ข้อมูลของท่านจะถูกเก็บเป็นความลับ และนำไปวิเคราะห์และเสนอผลงานในภาพรวมและใช้ประโยชน์สำหรับการศึกษาเท่านั้น

หากท่านมีข้อสงสัยใดๆ ดิฉันยินดีตอบข้อสงสัยของท่าน และขอขอบพระคุณทุกท่านที่ให้ความร่วมมือในการศึกษาครั้งนี้

สามารถติดต่อดิฉันได้ที่ โรงพยาบาลจิตเวชสงขลาราชนครินทร์ อำเภอเมือง จังหวัดสงขลา หรือโทรศัพท์หมายเลข 087-8377216 ได้ตลอดเวลา

นางสุดา ยุทธโท
นักศึกษาปริญญาโท สาธารณสุขศาสตรมหาบัณฑิต สาขาสาธารณสุขชุมชน
มหาวิทยาลัยราชภัฏสงขลา

SONGKHLA RAJABHAT UNIVERSITY

## แบบวัดสำหรับการวิจัย

**เรื่อง** ความสัมพันธ์ระหว่างความเชื่ออำนาจภายในตน-ภายนอกตนด้านสุขภาพกับภาวะซึมเศร้าในผู้ป่วยโรคซึมเศร้า

**คำชี้แจง** แบบวัดนี้ใช้สอบถามผู้ป่วยโรคซึมเศร้าที่มารับบริการที่โรงพยาบาลจิตเวชสงขลาราชนครินทร์ แผนกผู้ป่วยนอก โดยแบ่งเป็น 2 ตอน ดังนี้

ตอนที่1 แบบสอบถามข้อมูลส่วนบุคคล จำนวน 8 ข้อ

ตอนที่2 แบบวัดความเชื่ออำนาจภายในตน-ภายนอกตนด้านสุขภาพ จำนวน 15 ข้อ

**ตอนที่1 แบบสอบถามข้อมูลส่วนบุคคล**

**คำชี้แจง** โปรดเขียนเครื่องหมาย ลงในช่องว่างให้ตรงกับความเป็นจริงที่สุด

1. เพศ
   ( ) ชาย ( ) หญิง
2. อายุ.....ปี
3. ศาสนา
   ( ) พุทธ ( ) คริสต์ ( ) อิสลาม ( ) อื่นๆ ระบุ.....
4. อาชีพ
   ( ) เกษตรกรรม ( ) รับจ้าง ( ) ค้าขาย ( ) ข้าราชการ
   ( ) ไม่ได้ประกอบอาชีพ ( ) อื่นๆ ระบุ.....
5. ระดับการศึกษา
   ( ) ไม่ได้รับการศึกษา ( ) ประถมศึกษา ( ) มัธยมศึกษา
   ( ) ต่ำกว่าปริญญาตรี ( ) ปวช. /ปวส ( ) ปริญญาตรี
   ( ) ปริญญาโท ( ) อื่นๆ ระบุ.....
6. รายได้ของครอบครัวต่อเดือน
   ( ) พอใช้ ( ) ไม่พอใช้ ( ) มีเงินเก็บ
7. ความสัมพันธ์ของบุคคลในครอบครัว
   ( ) มีสัมพันธภาพกันดี ( ) มีสัมพันธภาพกันไม่ดี
8. การอบรมเลี้ยงดูที่ผ่านมา
   ( ) แบบเข้มงวด ( ) แบบปล่อยปละละเลย ( ) แบบอิสระเสรี

**ตอนที่ 2 แบบวัดความเชื่ออำนาจภายในตน-ภายนอกตนด้านสุขภาพ**

**คำชี้แจง**

1. โปรดเขียนเครื่องหมาย/ลงในช่องว่างให้ตรงกับความรู้สึกอย่างแท้จริง โดยตอบข้อละ 1 ช่อง เท่านั้น

2. ในกรณีท่านเห็นด้วย หมายถึง ท่านเห็นว่าสถานการณ์ในข้อนั้นตรงกับความรู้สึก โดยแบ่งเป็น

2.1 เห็นด้วยน้อย หมายถึง สถานการณ์ในข้อนั้นตรงกับความรู้สึกน้อย/ไม่มีผลต่อความรู้สึกหรือการดำเนินกิจวัตรประจำวัน

2.2 เห็นด้วยปานกลาง หมายถึง สถานการณ์ในข้อนั้นตรงกับความรู้สึกปานกลาง/ไม่มากหรือน้อย

2.3 เห็นด้วยมาก หมายถึง สถานการณ์ในข้อนั้นตรงกับความรู้สึกมาก/มีผลต่ออารมณ์ ความรู้สึกมากหรือมีผลต่อการดำเนินกิจวัตรประจำวัน

3. ในกรณีท่านไม่เห็นด้วย หมายถึง ท่านเห็นว่าสถานการณ์ในข้อนั้นตรงกับความรู้สึก โดยแบ่งเป็น

3.1 ไม่เห็นด้วยน้อย หมายถึง สถานการณ์ในข้อนั้นไม่ตรงกับความรู้สึก

3.2 ไม่เห็นด้วยปานกลาง หมายถึง สถานการณ์ในข้อนั้นไม่ตรงกับความรู้สึกปานกลาง

3.3 ไม่เห็นด้วยมาก หมายถึงสถานการณ์ในข้อนั้นไม่ตรงกับความรู้สึกมาก

SONGKHLA RAJABHAT UNIVERSITY

| เห็นด้วย | | | สถานการณ์ | ไม่เห็นด้วย | | |
|---|---|---|---|---|---|---|
| น้อย | ปานกลาง | มาก | | มาก | ปานกลาง | น้อย |
| | | | 1. เมื่อท่านรู้สึกเบื่อ ซึม หมดกำลังใจ และท้อแท้จนไม่อยากทำอะไร ท่านจะเป็นผู้กำหนดเองว่าท่านจะมีอาการดีขึ้นกลับมาเหมือนเดิมได้ในไม่ช้า | | | |
| | | | 2. วิธีที่ดีที่สุดสำหรับท่านที่จะเป็นช่องทางไม่ให้เกิดอาการซึมเศร้าคือการไปพบแพทย์อย่างสม่ำเสมอ | | | |
| | | | 3. สิ่งที่ส่งผลกระทบทำให้ท่านมีความรู้สึกเศร้า ท้อแท้ หมดหวัง หมดกำลังใจ ส่วนใหญ่แล้วมักเกิดขึ้นโดยความบังเอิญ | | | |
| | | | 4. ท่านสามารถดูแลตนเองไม่ให้เกิดความรู้สึกเศร้า | | | |
| | | | 5. บุคคลในครอบครัว ญาติพี่น้องและเพื่อนของท่านมีส่วนอย่างมากที่จะทำให้ท่านเกิดอาการเบื่อหน่าย เศร้า ท้อแท้ | | | |
| | | | 6. ไม่ว่าท่านจะทำตัวอย่างไรก็ตาม บางโอกาสท่านจะรู้สึกมีความรู้สึกเศร้า เสียใจขึ้นได้ | | | |
| | | | 7. ท่านมีความรู้สึกเป็นสุข สบายใจ เนื่องมาจากท่านปฏิบัติตัวได้ดี | | | |
| | | | 8. เมื่อท่านมีความรู้สึกเศร้าเกิดขึ้นแล้ว ท่านจะไปปรึกษาแพทย์หรือบุคลากรทางด้านสาธารณสุข | | | |
| | | | 9. สิ่งที่นับว่าโชคดีมากคือการที่ท่านมีสุขภาพจิตดีและมีจิตใจเข้มแข็งไม่มีอาการเศร้า หมดหวัง รู้สึกว่าตนเองไร้ค่า | | | |
| | | | 10. สิ่งสำคัญอย่างหนึ่งที่มีผลทำให้ไม่เกิดความรู้สึกหว้าเหว่เหงาไร้ค่า คือการปฏิบัติของท่านเอง | | | |
| | | | 11. การที่ท่านหายจากอาการเศร้า หมดกำลังใจ คิดว่าตนเองไม่มีคุณค่าได้นั้นเป็นเพราะแพทย์ พยาบาล ญาติพี่น้องในครอบครัว และเพื่อน เป็นส่วนหนึ่งในการดูแลท่านเป็นอย่างดี | | | |
| | | | 12. ถึงแม้ว่าท่านจะดูแลสุขภาพเป็นอย่างดี แต่ท่านก็ยังมีโอกาสเกิดอาการเศร้า ท้อแท้ได้ | | | |
| | | | 13. เมื่อท่านรู้สึกเศร้า เสียใจ ท่านมักคิดว่าตัวท่านเองเป็นต้นเหตุให้ท่านเกิดความรู้สึกนี้ | | | |
| | | | 14. ถ้าท่านปฏิบัติตามคำแนะนำของแพทย์เจ้าของไข้ของท่านอย่างเคร่งครัด จะส่งผลให้ท่านมีความรู้สึกเป็นสุขสบายใจ | | | |
| | | | 15. ความรู้สึกเศร้า เบื่อหน่าย รู้สึกเหนื่อยง่าย ไม่อยากทำอะไร สามารถเกิดขึ้นกับทุกคน โดยไม่สามารถหลีกเลี่ยงได้ | | | |

# ภาคผนวก ฉ

## สถิติที่ใช้ในการวิเคราะห์ข้อมูล

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

## สถิติที่ใช้ในการวิจัย

1. ค่า IOC แทน ดัชนีความสอดคล้องระหว่างข้อคำถาม

$$IOC = \frac{\sum R}{N}$$

เมื่อ IOC แทน ดัชนีความสอดคล้องระหว่างข้อคำถามตามวัตถุประสงค์ที่ได้ระบุไว้

$\Sigma^R$ แทน ผลรวมของคะแนนความคิดเห็นของผู้เชี่ยวชาญเนื้อหาทั้งหมด

N แทน จำนวนผู้เชี่ยวชาญ

2. ค่าเฉลี่ย (Mean)

$$\overline{X} = \frac{\sum X}{n}$$

เมื่อ $\overline{X}$ แทน ค่าเฉลี่ย

$\Sigma^X$ แทน ผลรวมของคะแนนทั้งหมดของกลุ่ม

n แทน จำนวนของคะแนนในกลุ่ม

3. ส่วนเบี่ยงเบนมาตรฐาน (Standard Deviation)

$$S.D. = \sqrt{\frac{n\sum X^2 - \left(\sum X\right)^2}{n(n-1)}}$$

เมื่อ S.D. แทน ค่าส่วนเบี่ยงเบนมาตรฐาน

X แทน ค่าคะแนน

n แทน จำนวนคะแนนในแต่ละกลุ่ม

$\Sigma$ แทน ผลรวม

4. ค่าความถี่ที่คาดว่าจะได้ $e_{ij}$ และค่าไคสแควร์

$$X^2_{calculated} = \sum_{t=1}^{r} \sum_{j=1}^{c} \frac{\left(o_{ij} - e_{ij}\right)^2}{e_{ij}}$$

# ภาคผนวก ช

## การวิเคราะห์ข้อมูลของกลุ่มตัวอย่าง

มหาวิทยาลัยราชภัฏสงขลา
SONGKHLA RAJABHAT UNIVERSITY

# Crosstabs

**Case Processing Summary**

| | Cases | | | | | |
|---|---|---|---|---|---|---|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| เพศ * GTO1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| เพศ * GTO2 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| เพศ * GTO3 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| เพศ * GTOTAL1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |

## เพศ * GTO1

**Crosstab**

| | | | GTO1 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| เพศ | ชาย | Count | 29 | 55 | 84 |
| | | % within เพศ | 34.5% | 65.5% | 100.0% |
| | | % within GTO1 | 35.4% | 47.0% | 42.2% |
| | หญิง | Count | 53 | 62 | 115 |
| | | % within เพศ | 46.1% | 53.9% | 100.0% |
| | | % within GTO1 | 64.6% | 53.0% | 57.8% |
| Total | | Count | 82 | 117 | 199 |
| | | % within เพศ | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | 2.679(b) | 1 | .102 | | |
| Continuity Correction(a) | 2.223 | 1 | .136 | | |
| Likelihood Ratio | 2.698 | 1 | .100 | | |
| Fisher's Exact Test | | | | .111 | .068 |
| Linear-by-Linear Association | 2.666 | 1 | .103 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 34.61.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | -.116 | .102 |
| | Cramer's V | .116 | .102 |
| | Contingency Coefficient | .115 | .102 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## เพศ * GTO2

**Crosstab**

| | | | GTO2 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| เพศ | ชาย | Count | 44 | 40 | 84 |
| | | % within เพศ | 52.4% | 47.6% | 100.0% |
| | | % within GTO2 | 50.0% | 36.0% | 42.2% |
| | หญิง | Count | 44 | 71 | 115 |
| | | % within เพศ | 38.3% | 61.7% | 100.0% |
| | | % within GTO2 | 50.0% | 64.0% | 57.8% |
| Total | | Count | 88 | 111 | 199 |
| | | % within เพศ | 44.2% | 55.8% | 100.0% |
| | | % within GTO2 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | 3.924(b) | 1 | .048 | | |
| Continuity Correction(a) | 3.372 | 1 | .066 | | |
| Likelihood Ratio | 3.925 | 1 | .048 | | |
| Fisher's Exact Test | | | | .060 | .033 |
| Linear-by-Linear Association | 3.904 | 1 | .048 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 37.15.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .140 | .048 |
| | Cramer's V | .140 | .048 |
| | Contingency Coefficient | .139 | .048 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## เพศ * GTO3

**Crosstab**

| | | | GTO3 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| เพศ | ชาย | Count | 43 | 41 | 84 |
| | | % within เพศ | 51.2% | 48.8% | 100.0% |
| | | % within GTO3 | 47.8% | 37.6% | 42.2% |
| | หญิง | Count | 47 | 68 | 115 |
| | | % within เพศ | 40.9% | 59.1% | 100.0% |
| | | % within GTO3 | 52.2% | 62.4% | 57.8% |
| Total | | Count | 90 | 109 | 199 |
| | | % within เพศ | 45.2% | 54.8% | 100.0% |
| | | % within GTO3 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | 2.087(b) | 1 | .149 | | |
| Continuity Correction(a) | 1.692 | 1 | .193 | | |
| Likelihood Ratio | 2.087 | 1 | .149 | | |
| Fisher's Exact Test | | | | .153 | .097 |
| Linear-by-Linear Association | 2.077 | 1 | .150 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 37.99.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .102 | .149 |
| | Cramer's V | .102 | .149 |
| | Contingency Coefficient | .102 | .149 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# เพศ * GTOTAL1

**Crosstab**

| | | | GTOTAL1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| เพศ | ชาย | Count | 36 | 48 | 84 |
| | | % within เพศ | 42.9% | 57.1% | 100.0% |
| | | % within GTOTAL1 | 43.4% | 41.4% | 42.2% |
| | หญิง | Count | 47 | 68 | 115 |
| | | % within เพศ | 40.9% | 59.1% | 100.0% |
| | | % within GTOTAL1 | 56.6% | 58.6% | 57.8% |
| Total | | Count | 83 | 116 | 199 |
| | | % within เพศ | 41.7% | 58.3% | 100.0% |
| | | % within GTOTAL1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | .079(b) | 1 | .779 | | |
| Continuity Correction(a) | .018 | 1 | .892 | | |
| Likelihood Ratio | .079 | 1 | .779 | | |
| Fisher's Exact Test | | | | .884 | .446 |
| Linear-by-Linear Association | .078 | 1 | .779 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 35.04.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .020 | .779 |
| | Cramer's V | .020 | .779 |
| | Contingency Coefficient | .020 | .779 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# Crosstabs

**Case Processing Summary**

| | Cases | | | | | |
|---|---|---|---|---|---|---|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| อายุ * GTO1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| อายุ * GTO2 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| อายุ * GTO3 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| อายุ * GTOTAL1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |

## อายุ * GTO1

**Crosstab**

| | | | GTO1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| อายุ | 19-29 ปี | Count | 14 | 24 | 38 |
| | | % within อายุ | 36.8% | 63.2% | 100.0% |
| | | % within GTO1 | 17.1% | 20.5% | 19.1% |
| | 30-39 ปี | Count | 16 | 28 | 44 |
| | | % within อายุ | 36.4% | 63.6% | 100.0% |
| | | % within GTO1 | 19.5% | 23.9% | 22.1% |
| | 40-49 ปี | Count | 25 | 27 | 52 |
| | | % within อายุ | 48.1% | 51.9% | 100.0% |
| | | % within GTO1 | 30.5% | 23.1% | 26.1% |
| | 50-59 ปี | Count | 27 | 38 | 65 |
| | | % within อายุ | 41.5% | 58.5% | 100.0% |
| | | % within GTO1 | 32.9% | 32.5% | 32.7% |
| Total | | Count | 82 | 117 | 199 |
| | | % within อายุ | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 1.741(a) | 3 | .628 |
| Likelihood Ratio | 1.737 | 3 | .629 |
| Linear-by-Linear Association | .536 | 1 | .464 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 15.66.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .094 | .628 |
| | Cramer's V | .094 | .628 |
| | Contingency Coefficient | .093 | .628 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# อายุ * GTO2

**Crosstab**

| | | | GTO2 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| อายุ | 19-29 ปี | Count | 22 | 16 | 38 |
| | | % within อายุ | 57.9% | 42.1% | 100.0% |
| | | % within GTO2 | 25.0% | 14.4% | 19.1% |
| | 30-39 ปี | Count | 19 | 25 | 44 |
| | | % within อายุ | 43.2% | 56.8% | 100.0% |
| | | % within GTO2 | 21.6% | 22.5% | 22.1% |
| | 40-49 ปี | Count | 25 | 27 | 52 |
| | | % within อายุ | 48.1% | 51.9% | 100.0% |
| | | % within GTO2 | 28.4% | 24.3% | 26.1% |
| | 50-59 ปี | Count | 22 | 43 | 65 |
| | | % within อายุ | 33.8% | 66.2% | 100.0% |
| | | % within GTO2 | 25.0% | 38.7% | 32.7% |
| Total | | Count | 88 | 111 | 199 |
| | | % within อายุ | 44.2% | 55.8% | 100.0% |
| | | % within GTO2 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 6.050(a) | 3 | .109 |
| Likelihood Ratio | 6.093 | 3 | .107 |
| Linear-by-Linear Association | 4.565 | 1 | .033 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 16.80.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .174 | .109 |
| | Cramer's V | .174 | .109 |
| | Contingency Coefficient | .172 | .109 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## อายุ * GTO3 **Crosstab**

| | | | GTO3 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| อายุ | 19-29 ปี | Count | 18 | 20 | 38 |
| | | % within อายุ | 47.4% | 52.6% | 100.0% |
| | | % within GTO3 | 20.0% | 18.3% | 19.1% |
| | 30-39 ปี | Count | 16 | 28 | 44 |
| | | % within อายุ | 36.4% | 63.6% | 100.0% |
| | | % within GTO3 | 17.8% | 25.7% | 22.1% |
| | 40-49 ปี | Count | 27 | 25 | 52 |
| | | % within อายุ | 51.9% | 48.1% | 100.0% |
| | | % within GTO3 | 30.0% | 22.9% | 26.1% |
| | 50-59 ปี | Count | 29 | 36 | 65 |
| | | % within อายุ | 44.6% | 55.4% | 100.0% |
| | | % within GTO3 | 32.2% | 33.0% | 32.7% |
| Total | | Count | 90 | 109 | 199 |
| | | % within อายุ | 45.2% | 54.8% | 100.0% |
| | | % within GTO3 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 2.417(a) | 3 | .491 |
| Likelihood Ratio | 2.435 | 3 | .487 |
| Linear-by-Linear Association | .057 | 1 | .811 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 17.19.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .110 | .491 |
| | Cramer's V | .110 | .491 |
| | Contingency Coefficient | .110 | .491 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# อายุ * GTOTAL1

**Crosstab**

| | | | GTOTAL1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| อายุ | 19-29 ปี | Count | 17 | 21 | 38 |
| | | % within อายุ | 44.7% | 55.3% | 100.0% |
| | | % within GTOTAL1 | 20.5% | 18.1% | 19.1% |
| | 30-39 ปี | Count | 16 | 28 | 44 |
| | | % within อายุ | 36.4% | 63.6% | 100.0% |
| | | % within GTOTAL1 | 19.3% | 24.1% | 22.1% |
| | 40-49 ปี | Count | 26 | 26 | 52 |
| | | % within อายุ | 50.0% | 50.0% | 100.0% |
| | | % within GTOTAL1 | 31.3% | 22.4% | 26.1% |
| | 50-59 ปี | Count | 24 | 41 | 65 |
| | | % within อายุ | 36.9% | 63.1% | 100.0% |
| | | % within GTOTAL1 | 28.9% | 35.3% | 32.7% |
| Total | | Count | 83 | 116 | 199 |
| | | % within อายุ | 41.7% | 58.3% | 100.0% |
| | | % within GTOTAL1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 2.743(a) | 3 | .433 |
| Likelihood Ratio | 2.737 | 3 | .434 |
| Linear-by-Linear Association | .156 | 1 | .693 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 15.85.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .117 | .433 |
| | Cramer's V | .117 | .433 |
| | Contingency Coefficient | .117 | .433 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# Crosstabs

**Case Processing Summary**

| | Cases | | | | | |
|---|---|---|---|---|---|---|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| ศาสนา * GTO1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| ศาสนา * GTO2 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| ศาสนา * GTO3 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| ศาสนา * GTOTAL1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |

## ศาสนา * GTO1 Crosstab

| | | | GTO1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| ศาสนา | พุทธ | Count | 65 | 88 | 153 |
| | | % within ศาสนา | 42.5% | 57.5% | 100.0% |
| | | % within GTO1 | 79.3% | 75.2% | 76.9% |
| | อื่นๆ | Count | 17 | 29 | 46 |
| | | % within ศาสนา | 37.0% | 63.0% | 100.0% |
| | | % within GTO1 | 20.7% | 24.8% | 23.1% |
| Total | | Count | 82 | 117 | 199 |
| | | % within ศาสนา | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | .446(b) | 1 | .504 | | |
| Continuity Correction(a) | .247 | 1 | .619 | | |
| Likelihood Ratio | .450 | 1 | .502 | | |
| Fisher's Exact Test | | | | .609 | .311 |
| Linear-by-Linear Association | .444 | 1 | .505 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 18.95.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .047 | .504 |
| | Cramer's V | .047 | .504 |
| | Contingency Coefficient | .047 | .504 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## ศาสนา * GTO2

**Crosstab**

| | | | GTO2 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| ศาสนา | พุทธ | Count | 73 | 80 | 153 |
| | | % within ศาสนา | 47.7% | 52.3% | 100.0% |
| | | % within GTO2 | 83.0% | 72.1% | 76.9% |
| | อื่นๆ | Count | 15 | 31 | 46 |
| | | % within ศาสนา | 32.6% | 67.4% | 100.0% |
| | | % within GTO2 | 17.0% | 27.9% | 23.1% |
| Total | | Count | 88 | 111 | 199 |
| | | % within ศาสนา | 44.2% | 55.8% | 100.0% |
| | | % within GTO2 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | 3.271(b) | 1 | .071 | | |
| Continuity Correction(a) | 2.687 | 1 | .101 | | |
| Likelihood Ratio | 3.339 | 1 | .068 | | |
| Fisher's Exact Test | | | | .090 | .050 |
| Linear-by-Linear Association | 3.254 | 1 | .071 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 20.34.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .128 | .071 |
| | Cramer's V | .128 | .071 |
| | Contingency Coefficient | .127 | .071 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## ศาสนา * GTO3

**Crosstab**

| | | | GTO3 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| ศาสนา | พุทธ | Count | 70 | 83 | 153 |
| | | % within ศาสนา | 45.8% | 54.2% | 100.0% |
| | | % within GTO3 | 77.8% | 76.1% | 76.9% |
| | อื่นๆ | Count | 20 | 26 | 46 |
| | | % within ศาสนา | 43.5% | 56.5% | 100.0% |
| | | % within GTO3 | 22.2% | 23.9% | 23.1% |
| Total | | Count | 90 | 109 | 199 |
| | | % within ศาสนา | 45.2% | 54.8% | 100.0% |
| | | % within GTO3 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | .074(b) | 1 | .786 | | |
| Continuity Correction(a) | .011 | 1 | .918 | | |
| Likelihood Ratio | .074 | 1 | .786 | | |
| Fisher's Exact Test | | | | .866 | .460 |
| Linear-by-Linear Association | .073 | 1 | .786 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 20.80.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .019 | .786 |
| | Cramer's V | .019 | .786 |
| | Contingency Coefficient | .019 | .786 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# ศาสนา * GTOTAL1

**Crosstab**

| | | | GTOTAL1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| ศาสนา | พุทธ | Count | 67 | 86 | 153 |
| | | % within ศาสนา | 43.8% | 56.2% | 100.0% |
| | | % within GTOTAL1 | 80.7% | 74.1% | 76.9% |
| | อื่นๆ | Count | 16 | 30 | 46 |
| | | % within ศาสนา | 34.8% | 65.2% | 100.0% |
| | | % within GTOTAL1 | 19.3% | 25.9% | 23.1% |
| Total | | Count | 83 | 116 | 199 |
| | | % within ศาสนา | 41.7% | 58.3% | 100.0% |
| | | % within GTOTAL1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | 1.180(b) | 1 | .277 | | |
| Continuity Correction(a) | .839 | 1 | .360 | | |
| Likelihood Ratio | 1.197 | 1 | .274 | | |
| Fisher's Exact Test | | | | .310 | .180 |
| Linear-by-Linear Association | 1.175 | 1 | .278 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 19.19.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .077 | .277 |
| | Cramer's V | .077 | .277 |
| | Contingency Coefficient | .077 | .277 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# Crosstabs

**Case Processing Summary**

| | Cases | | | | | |
|---|---|---|---|---|---|---|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| อาชีพ * GTO1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| อาชีพ * GTO2 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| อาชีพ * GTO3 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| อาชีพ * GTOTAL1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |

# อาชีพ * GTO1

**Crosstab**

| | | | GTO1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| อาชีพ | เกษตรกร | Count | 34 | 43 | 77 |
| | | % within อาชีพ | 44.2% | 55.8% | 100.0% |
| | | % within GTO1 | 41.5% | 36.8% | 38.7% |
| | รับจ้าง | Count | 13 | 25 | 38 |
| | | % within อาชีพ | 34.2% | 65.8% | 100.0% |
| | | % within GTO1 | 15.9% | 21.4% | 19.1% |
| | ธุรกิจส่วนตัว | Count | 8 | 17 | 25 |
| | | % within อาชีพ | 32.0% | 68.0% | 100.0% |
| | | % within GTO1 | 9.8% | 14.5% | 12.6% |
| | รับราชการ | Count | 16 | 17 | 33 |
| | | % within อาชีพ | 48.5% | 51.5% | 100.0% |
| | | % within GTO1 | 19.5% | 14.5% | 16.6% |
| | ไม่ได้ประกอบอาชีพ | Count | 11 | 15 | 26 |
| | | % within อาชีพ | 42.3% | 57.7% | 100.0% |
| | | % within GTO1 | 13.4% | 12.8% | 13.1% |
| Total | | Count | 82 | 117 | 199 |
| | | % within อาชีพ | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 2.653(a) | 4 | .617 |
| Likelihood Ratio | 2.684 | 4 | .612 |
| Linear-by-Linear Association | .011 | 1 | .915 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 10.30.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .115 | .617 |
| | Cramer's V | .115 | .617 |
| | Contingency Coefficient | .115 | .617 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# อาชีพ * GTO2

**Crosstab**

| | | | GTO2 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| อาชีพ | เกษตรกร | Count | 28 | 49 | 77 |
| | | % within อาชีพ | 36.4% | 63.6% | 100.0% |
| | | % within GTO2 | 31.8% | 44.1% | 38.7% |
| | รับจ้าง | Count | 16 | 22 | 38 |
| | | % within อาชีพ | 42.1% | 57.9% | 100.0% |
| | | % within GTO2 | 18.2% | 19.8% | 19.1% |
| | ธุรกิจส่วนตัว | Count | 9 | 16 | 25 |
| | | % within อาชีพ | 36.0% | 64.0% | 100.0% |
| | | % within GTO2 | 10.2% | 14.4% | 12.6% |
| | รับราชการ | Count | 17 | 16 | 33 |
| | | % within อาชีพ | 51.5% | 48.5% | 100.0% |
| | | % within GTO2 | 19.3% | 14.4% | 16.6% |
| | ไม่ได้ประกอบอาชีพ | Count | 18 | 8 | 26 |
| | | % within อาชีพ | 69.2% | 30.8% | 100.0% |
| | | % within GTO2 | 20.5% | 7.2% | 13.1% |
| Total | | Count | 88 | 111 | 199 |
| | | % within อาชีพ | 44.2% | 55.8% | 100.0% |
| | | % within GTO2 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 9.986(a) | 4 | .041 |
| Likelihood Ratio | 10.051 | 4 | .040 |
| Linear-by-Linear Association | 7.602 | 1 | .006 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.06.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .224 | .041 |
| | Cramer's V | .224 | .041 |
| | Contingency Coefficient | .219 | .041 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## อาชีพ * GTO3 Crosstab

| | | | GTO3 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| อาชีพ | เกษตรกร | Count | 31 | 46 | 77 |
| | | % within อาชีพ | 40.3% | 59.7% | 100.0% |
| | | % within GTO3 | 34.4% | 42.2% | 38.7% |
| | รับจ้าง | Count | 14 | 24 | 38 |
| | | % within อาชีพ | 36.8% | 63.2% | 100.0% |
| | | % within GTO3 | 15.6% | 22.0% | 19.1% |
| | ธุรกิจส่วนตัว | Count | 13 | 12 | 25 |
| | | % within อาชีพ | 52.0% | 48.0% | 100.0% |
| | | % within GTO3 | 14.4% | 11.0% | 12.6% |
| | รับราชการ | Count | 18 | 15 | 33 |
| | | % within อาชีพ | 54.5% | 45.5% | 100.0% |
| | | % within GTO3 | 20.0% | 13.8% | 16.6% |
| | ไม่ได้ประกอบอาชีพ | Count | 14 | 12 | 26 |
| | | % within อาชีพ | 53.8% | 46.2% | 100.0% |
| | | % within GTO3 | 15.6% | 11.0% | 13.1% |
| Total | | Count | 90 | 109 | 199 |
| | | % within อาชีพ | 45.2% | 54.8% | 100.0% |
| | | % within GTO3 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 4.245(a) | 4 | .374 |
| Likelihood Ratio | 4.254 | 4 | .373 |
| Linear-by-Linear Association | 3.193 | 1 | .074 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.31.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .146 | .374 |
| | Cramer's V | .146 | .374 |
| | Contingency Coefficient | .145 | .374 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# อาชีพ * GTOTAL1

**Crosstab**

| | | | GTOTAL1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| อาชีพ | เกษตรกร | Count | 27 | 50 | 77 |
| | | % within อาชีพ | 35.1% | 64.9% | 100.0% |
| | | % within GTOTAL1 | 32.5% | 43.1% | 38.7% |
| | รับจ้าง | Count | 15 | 23 | 38 |
| | | % within อาชีพ | 39.5% | 60.5% | 100.0% |
| | | % within GTOTAL1 | 18.1% | 19.8% | 19.1% |
| | ธุรกิจส่วนตัว | Count | 9 | 16 | 25 |
| | | % within อาชีพ | 36.0% | 64.0% | 100.0% |
| | | % within GTOTAL1 | 10.8% | 13.8% | 12.6% |
| | รับราชการ | Count | 17 | 16 | 33 |
| | | % within อาชีพ | 51.5% | 48.5% | 100.0% |
| | | % within GTOTAL1 | 20.5% | 13.8% | 16.6% |
| | ไม่ได้ประกอบอาชีพ | Count | 15 | 11 | 26 |
| | | % within อาชีพ | 57.7% | 42.3% | 100.0% |
| | | % within GTOTAL1 | 18.1% | 9.5% | 13.1% |
| Total | | Count | 83 | 116 | 199 |
| | | % within อาชีพ | 41.7% | 58.3% | 100.0% |
| | | % within GTOTAL1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 5.848(a) | 4 | .211 |
| Likelihood Ratio | 5.810 | 4 | .214 |
| Linear-by-Linear Association | 4.925 | 1 | .026 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 10.43.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .171 | .211 |
| | Cramer's V | .171 | .211 |
| | Contingency Coefficient | .169 | .211 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# Crosstabs

**Case Processing Summary**

| | Cases | | | | | |
|---|---|---|---|---|---|---|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| การศึกษา * GTO1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| การศึกษา * GTO2 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| การศึกษา * GTO3 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| การศึกษา * GTOTAL1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |

# การศึกษา * GTO1

**Crosstab**

| | | | GTO1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| การศึกษา | ประถมศึกษา | Count | 39 | 46 | 85 |
| | | % within การศึกษา | 45.9% | 54.1% | 100.0% |
| | | % within GTO1 | 47.6% | 39.3% | 42.7% |
| | มัธยมศึกษา | Count | 18 | 32 | 50 |
| | | % within การศึกษา | 36.0% | 64.0% | 100.0% |
| | | % within GTO1 | 22.0% | 27.4% | 25.1% |
| | อนุปริญญา/เทียบเท่า | Count | 7 | 18 | 25 |
| | | % within การศึกษา | 28.0% | 72.0% | 100.0% |
| | | % within GTO1 | 8.5% | 15.4% | 12.6% |
| | ปริญญาตรีและสูงกว่า | Count | 18 | 21 | 39 |
| | | % within การศึกษา | 46.2% | 53.8% | 100.0% |
| | | % within GTO1 | 22.0% | 17.9% | 19.6% |
| Total | | Count | 82 | 117 | 199 |
| | | % within การศึกษา | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 3.520(a) | 3 | .318 |
| Likelihood Ratio | 3.603 | 3 | .308 |
| Linear-by-Linear Association | .181 | 1 | .670 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 10.30.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .133 | .318 |
| | Cramer's V | .133 | .318 |
| | Contingency Coefficient | .132 | .318 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# การศึกษา * GTO2

**Crosstab**

| | | | GTO2 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| การศึกษา | ประถมศึกษา | Count | 38 | 47 | 85 |
| | | % within การศึกษา | 44.7% | 55.3% | 100.0% |
| | | % within GTO2 | 43.2% | 42.3% | 42.7% |
| | มัธยมศึกษา | Count | 20 | 30 | 50 |
| | | % within การศึกษา | 40.0% | 60.0% | 100.0% |
| | | % within GTO2 | 22.7% | 27.0% | 25.1% |
| | อนุปริญญา/เทียบเท่า | Count | 12 | 13 | 25 |
| | | % within การศึกษา | 48.0% | 52.0% | 100.0% |
| | | % within GTO2 | 13.6% | 11.7% | 12.6% |
| | ปริญญาตรีและสูงกว่า | Count | 18 | 21 | 39 |
| | | % within การศึกษา | 46.2% | 53.8% | 100.0% |
| | | % within GTO2 | 20.5% | 18.9% | 19.6% |
| Total | | Count | 88 | 111 | 199 |
| | | % within การศึกษา | 44.2% | 55.8% | 100.0% |
| | | % within GTO2 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | .573(a) | 3 | .903 |
| Likelihood Ratio | .575 | 3 | .902 |
| Linear-by-Linear Association | .064 | 1 | .801 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.06.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .054 | .903 |
| | Cramer's V | .054 | .903 |
| | Contingency Coefficient | .054 | .903 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# การศึกษา * GTO3

**Crosstab**

| | | | GTO3 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| การศึกษา | ประถมศึกษา | Count | 36 | 49 | 85 |
| | | % within การศึกษา | 42.4% | 57.6% | 100.0% |
| | | % within GTO3 | 40.0% | 45.0% | 42.7% |
| | มัธยมศึกษา | Count | 21 | 29 | 50 |
| | | % within การศึกษา | 42.0% | 58.0% | 100.0% |
| | | % within GTO3 | 23.3% | 26.6% | 25.1% |
| | อนุปริญญา/เทียบเท่า | Count | 13 | 12 | 25 |
| | | % within การศึกษา | 52.0% | 48.0% | 100.0% |
| | | % within GTO3 | 14.4% | 11.0% | 12.6% |
| | ปริญญาตรีและสูงกว่า | Count | 20 | 19 | 39 |
| | | % within การศึกษา | 51.3% | 48.7% | 100.0% |
| | | % within GTO3 | 22.2% | 17.4% | 19.6% |
| Total | | Count | 90 | 109 | 199 |
| | | % within การศึกษา | 45.2% | 54.8% | 100.0% |
| | | % within GTO3 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 1.534(a) | 3 | .674 |
| Likelihood Ratio | 1.530 | 3 | .675 |
| Linear-by-Linear Association | 1.192 | 1 | .275 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 11.31.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .088 | .674 |
| | Cramer's V | .088 | .674 |
| | Contingency Coefficient | .087 | .674 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# การศึกษา * GTOTAL1

**Crosstab**

| | | | GTOTAL1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| การศึกษา | ประถมศึกษา | Count | 35 | 50 | 85 |
| | | % within การศึกษา | 41.2% | 58.8% | 100.0% |
| | | % within GTOTAL1 | 42.2% | 43.1% | 42.7% |
| | มัธยมศึกษา | Count | 20 | 30 | 50 |
| | | % within การศึกษา | 40.0% | 60.0% | 100.0% |
| | | % within GTOTAL1 | 24.1% | 25.9% | 25.1% |
| | อนุปริญญา/เทียบเท่า | Count | 9 | 16 | 25 |
| | | % within การศึกษา | 36.0% | 64.0% | 100.0% |
| | | % within GTOTAL1 | 10.8% | 13.8% | 12.6% |
| | ปริญญาตรีและสูงกว่า | Count | 19 | 20 | 39 |
| | | % within การศึกษา | 48.7% | 51.3% | 100.0% |
| | | % within GTOTAL1 | 22.9% | 17.2% | 19.6% |
| Total | | Count | 83 | 116 | 199 |
| | | % within การศึกษา | 41.7% | 58.3% | 100.0% |
| | | % within GTOTAL1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 1.193(a) | 3 | .755 |
| Likelihood Ratio | 1.189 | 3 | .756 |
| Linear-by-Linear Association | .312 | 1 | .576 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 10.43.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .077 | .755 |
| | Cramer's V | .077 | .755 |
| | Contingency Coefficient | .077 | .755 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# Crosstabs

**Case Processing Summary**

| | Cases | | | | | |
|---|---|---|---|---|---|---|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| รายได้ * GTO1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| รายได้ * GTO2 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| รายได้ * GTO3 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| รายได้ * GTOTAL1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |

# รายได้ * GTO1

**Crosstab**

| | | | GTO1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| รายได้ | พอใช้ | Count | 63 | 100 | 163 |
| | | % within รายได้ | 38.7% | 61.3% | 100.0% |
| | | % within GTO1 | 76.8% | 85.5% | 81.9% |
| | ไม่พอใช้ | Count | 9 | 9 | 18 |
| | | % within รายได้ | 50.0% | 50.0% | 100.0% |
| | | % within GTO1 | 11.0% | 7.7% | 9.0% |
| | มีเงินเก็บ | Count | 10 | 8 | 18 |
| | | % within รายได้ | 55.6% | 44.4% | 100.0% |
| | | % within GTO1 | 12.2% | 6.8% | 9.0% |
| Total | | Count | 82 | 117 | 199 |
| | | % within รายได้ | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 2.544(a) | 2 | .280 |
| Likelihood Ratio | 2.507 | 2 | .285 |
| Linear-by-Linear Association | 2.483 | 1 | .115 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.42.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .113 | .280 |
| | Cramer's V | .113 | .280 |
| | Contingency Coefficient | .112 | .280 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# รายได้ * GTO2

**Crosstab**

| | | | GTO2 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| รายได้ | พอใช้ | Count | 67 | 96 | 163 |
| | | % within รายได้ | 41.1% | 58.9% | 100.0% |
| | | % within GTO2 | 76.1% | 86.5% | 81.9% |
| | ไม่พอใช้ | Count | 9 | 9 | 18 |
| | | % within รายได้ | 50.0% | 50.0% | 100.0% |
| | | % within GTO2 | 10.2% | 8.1% | 9.0% |
| | มีเงินเก็บ | Count | 12 | 6 | 18 |
| | | % within รายได้ | 66.7% | 33.3% | 100.0% |
| | | % within GTO2 | 13.6% | 5.4% | 9.0% |
| Total | | Count | 88 | 111 | 199 |
| | | % within รายได้ | 44.2% | 55.8% | 100.0% |
| | | % within GTO2 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 4.562(a) | 2 | .102 |
| Likelihood Ratio | 4.562 | 2 | .102 |
| Linear-by-Linear Association | 4.453 | 1 | .035 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.96.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .151 | .102 |
| | Cramer's V | .151 | .102 |
| | Contingency Coefficient | .150 | .102 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# รายได้ * GTO3

**Crosstab**

| | | | GTO3 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| รายได้ | พอใช้ | Count | 74 | 89 | 163 |
| | | % within รายได้ | 45.4% | 54.6% | 100.0% |
| | | % within GTO3 | 82.2% | 81.7% | 81.9% |
| | ไม่พอใช้ | Count | 7 | 11 | 18 |
| | | % within รายได้ | 38.9% | 61.1% | 100.0% |
| | | % within GTO3 | 7.8% | 10.1% | 9.0% |
| | มีเงินเก็บ | Count | 9 | 9 | 18 |
| | | % within รายได้ | 50.0% | 50.0% | 100.0% |
| | | % within GTO3 | 10.0% | 8.3% | 9.0% |
| Total | | Count | 90 | 109 | 199 |
| | | % within รายได้ | 45.2% | 54.8% | 100.0% |
| | | % within GTO3 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | .459(a) | 2 | .795 |
| Likelihood Ratio | .462 | 2 | .794 |
| Linear-by-Linear Association | .018 | 1 | .894 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 8.14.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .048 | .795 |
| | Cramer's V | .048 | .795 |
| | Contingency Coefficient | .048 | .795 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# รายได้ * GTOTAL1

**Crosstab**

| | | | GTOTAL1 | | |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | Total |
| รายได้ | พอใช้ | Count | 64 | 99 | 163 |
| | | % within รายได้ | 39.3% | 60.7% | 100.0% |
| | | % within GTOTAL1 | 77.1% | 85.3% | 81.9% |
| | ไม่พอใช้ | Count | 10 | 8 | 18 |
| | | % within รายได้ | 55.6% | 44.4% | 100.0% |
| | | % within GTOTAL1 | 12.0% | 6.9% | 9.0% |
| | มีเงินเก็บ | Count | 9 | 9 | 18 |
| | | % within รายได้ | 50.0% | 50.0% | 100.0% |
| | | % within GTOTAL1 | 10.8% | 7.8% | 9.0% |
| Total | | Count | 83 | 116 | 199 |
| | | % within รายได้ | 41.7% | 58.3% | 100.0% |
| | | % within GTOTAL1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 2.329(a) | 2 | .312 |
| Likelihood Ratio | 2.299 | 2 | .317 |
| Linear-by-Linear Association | 1.630 | 1 | .202 |
| N of Valid Cases | 199 | | |

a 0 cells (.0%) have expected count less than 5. The minimum expected count is 7.51.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .108 | .312 |
| | Cramer's V | .108 | .312 |
| | Contingency Coefficient | .108 | .312 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# Crosstabs

**Case Processing Summary**

| | Cases | | | | | |
|---|---|---|---|---|---|---|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| สัมพันธ์ * GTO1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| สัมพันธ์ * GTO2 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| สัมพันธ์ * GTO3 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| สัมพันธ์ * GTOTAL1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |

# สัมพันธ์ * GTO1

**Crosstab**

| | | | GTO1 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| สัมพันธ์ | มีความสัมพันธ์กันดี | Count | 74 | 110 | 184 |
| | | % within สัมพันธ์ | 40.2% | 59.8% | 100.0% |
| | | % within GTO1 | 90.2% | 94.0% | 92.5% |
| | มีความสัมพันธ์กันไม่ดี | Count | 8 | 7 | 15 |
| | | % within สัมพันธ์ | 53.3% | 46.7% | 100.0% |
| | | % within GTO1 | 9.8% | 6.0% | 7.5% |
| Total | | Count | 82 | 117 | 199 |
| | | % within สัมพันธ์ | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | .985(b) | 1 | .321 | | |
| Continuity Correction(a) | .518 | 1 | .472 | | |
| Likelihood Ratio | .968 | 1 | .325 | | |
| Fisher's Exact Test | | | | .415 | .234 |
| Linear-by-Linear Association | .980 | 1 | .322 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.18.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | -.070 | .321 |
| | Cramer's V | .070 | .321 |
| | Contingency Coefficient | .070 | .321 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## สัมพันธ์ * GTO2

**Crosstab**

| | | | GTO2 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| สัมพันธ์ | มีความสัมพันธ์กันดี | Count | 84 | 100 | 184 |
| | | % within สัมพันธ์ | 45.7% | 54.3% | 100.0% |
| | | % within GTO2 | 95.5% | 90.1% | 92.5% |
| | มีความสัมพันธ์กันไม่ดี | Count | 4 | 11 | 15 |
| | | % within สัมพันธ์ | 26.7% | 73.3% | 100.0% |
| | | % within GTO2 | 4.5% | 9.9% | 7.5% |
| Total | | Count | 88 | 111 | 199 |
| | | % within สัมพันธ์ | 44.2% | 55.8% | 100.0% |
| | | % within GTO2 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | 2.027(b) | 1 | .155 | | |
| Continuity Correction(a) | 1.330 | 1 | .249 | | |
| Likelihood Ratio | 2.126 | 1 | .145 | | |
| Fisher's Exact Test | | | | .184 | .123 |
| Linear-by-Linear Association | 2.017 | 1 | .156 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.63.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .101 | .155 |
| | Cramer's V | .101 | .155 |
| | Contingency Coefficient | .100 | .155 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## สัมพันธ์ * GTO3

**Crosstab**

| | | | GTO3 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| สัมพันธ์ | มีความสัมพันธ์กันดี | Count | 84 | 100 | 184 |
| | | % within สัมพันธ์ | 45.7% | 54.3% | 100.0% |
| | | % within GTO3 | 93.3% | 91.7% | 92.5% |
| | มีความสัมพันธ์กันไม่ดี | Count | 6 | 9 | 15 |
| | | % within สัมพันธ์ | 40.0% | 60.0% | 100.0% |
| | | % within GTO3 | 6.7% | 8.3% | 7.5% |
| Total | | Count | 90 | 109 | 199 |
| | | % within สัมพันธ์ | 45.2% | 54.8% | 100.0% |
| | | % within GTO3 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | .179(b) | 1 | .672 | | |
| Continuity Correction(a) | .023 | 1 | .878 | | |
| Likelihood Ratio | .180 | 1 | .671 | | |
| Fisher's Exact Test | | | | .790 | .442 |
| Linear-by-Linear Association | .178 | 1 | .673 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.78.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .030 | .672 |
| | Cramer's V | .030 | .672 |
| | Contingency Coefficient | .030 | .672 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## สัมพันธ์ * GTOTAL1

**Crosstab**

| | | | GTOTAL1 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| สัมพันธ์ | มีความสัมพันธ์กันดี | Count | 78 | 106 | 184 |
| | | % within สัมพันธ์ | 42.4% | 57.6% | 100.0% |
| | | % within GTOTAL1 | 94.0% | 91.4% | 92.5% |
| | มีความสัมพันธ์กันไม่ดี | Count | 5 | 10 | 15 |
| | | % within สัมพันธ์ | 33.3% | 66.7% | 100.0% |
| | | % within GTOTAL1 | 6.0% | 8.6% | 7.5% |
| Total | | Count | 83 | 116 | 199 |
| | | % within สัมพันธ์ | 41.7% | 58.3% | 100.0% |
| | | % within GTOTAL1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) | Exact Sig. (2-sided) | Exact Sig. (1-sided) |
|---|---|---|---|---|---|
| Pearson Chi-Square | .468(b) | 1 | .494 | | |
| Continuity Correction(a) | .170 | 1 | .680 | | |
| Likelihood Ratio | .479 | 1 | .489 | | |
| Fisher's Exact Test | | | | .592 | .345 |
| Linear-by-Linear Association | .466 | 1 | .495 | | |
| N of Valid Cases | 199 | | | | |

a Computed only for a 2x2 table
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.26.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .048 | .494 |
| | Cramer's V | .048 | .494 |
| | Contingency Coefficient | .048 | .494 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# Crosstabs

**Case Processing Summary**

| | Cases | | | | | |
|---|---|---|---|---|---|---|
| | Valid | | Missing | | Total | |
| | N | Percent | N | Percent | N | Percent |
| เลี้ยงดู * GTO1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| เลี้ยงดู * GTO2 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| เลี้ยงดู * GTO3 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |
| เลี้ยงดู * GTOTAL1 | 199 | 100.0% | 0 | .0% | 199 | 100.0% |

# เลี้ยงดู * GTO1

**Crosstab**

| | | | GTO1 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| เลี้ยงดู | แบบเข้มงวด | Count | 18 | 25 | 43 |
| | | % within เลี้ยงดู | 41.9% | 58.1% | 100.0% |
| | | % within GTO1 | 22.0% | 21.4% | 21.6% |
| | แบบปล่อยปละละเลย | Count | 3 | 5 | 8 |
| | | % within เลี้ยงดู | 37.5% | 62.5% | 100.0% |
| | | % within GTO1 | 3.7% | 4.3% | 4.0% |
| | แบบอิสระเสรี | Count | 61 | 87 | 148 |
| | | % within เลี้ยงดู | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 74.4% | 74.4% | 74.4% |
| Total | | Count | 82 | 117 | 199 |
| | | % within เลี้ยงดู | 41.2% | 58.8% | 100.0% |
| | | % within GTO1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | .053(a) | 2 | .974 |
| Likelihood Ratio | .053 | 2 | .974 |
| Linear-by-Linear Association | .002 | 1 | .963 |
| N of Valid Cases | 199 | | |

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 3.30.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .016 | .974 |
| | Cramer's V | .016 | .974 |
| | Contingency Coefficient | .016 | .974 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# เลี้ยงดู * GTO2

**Crosstab**

| | | | GTO2 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| เลี้ยงดู | แบบเข้มงวด | Count | 22 | 21 | 43 |
| | | % within เลี้ยงดู | 51.2% | 48.8% | 100.0% |
| | | % within GTO2 | 25.0% | 18.9% | 21.6% |
| | แบบปล่อยปละละเลย | Count | 3 | 5 | 8 |
| | | % within เลี้ยงดู | 37.5% | 62.5% | 100.0% |
| | | % within GTO2 | 3.4% | 4.5% | 4.0% |
| | แบบอิสระเสรี | Count | 63 | 85 | 148 |
| | | % within เลี้ยงดู | 42.6% | 57.4% | 100.0% |
| | | % within GTO2 | 71.6% | 76.6% | 74.4% |
| Total | | Count | 88 | 111 | 199 |
| | | % within เลี้ยงดู | 44.2% | 55.8% | 100.0% |
| | | % within GTO2 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 1.151(a) | 2 | .563 |
| Likelihood Ratio | 1.147 | 2 | .564 |
| Linear-by-Linear Association | .878 | 1 | .349 |
| N of Valid Cases | 199 | | |

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 3.54.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .076 | .563 |
| | Cramer's V | .076 | .563 |
| | Contingency Coefficient | .076 | .563 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

## เลี้ยงดู * GTO3

**Crosstab**

| | | | GTO3 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| เลี้ยงดู | แบบเข้มงวด | Count | 23 | 20 | 43 |
| | | % within เลี้ยงดู | 53.5% | 46.5% | 100.0% |
| | | % within GTO3 | 25.6% | 18.3% | 21.6% |
| | แบบปล่อยปละละเลย | Count | 2 | 6 | 8 |
| | | % within เลี้ยงดู | 25.0% | 75.0% | 100.0% |
| | | % within GTO3 | 2.2% | 5.5% | 4.0% |
| | แบบอิสระเสรี | Count | 65 | 83 | 148 |
| | | % within เลี้ยงดู | 43.9% | 56.1% | 100.0% |
| | | % within GTO3 | 72.2% | 76.1% | 74.4% |
| Total | | Count | 90 | 109 | 199 |
| | | % within เลี้ยงดู | 45.2% | 54.8% | 100.0% |
| | | % within GTO3 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | 2.608(a) | 2 | .271 |
| Likelihood Ratio | 2.680 | 2 | .262 |
| Linear-by-Linear Association | .892 | 1 | .345 |
| N of Valid Cases | 199 | | |

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 3.62.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .114 | .271 |
| | Cramer's V | .114 | .271 |
| | Contingency Coefficient | .114 | .271 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.

# เลี้ยงดู * GTOTAL1

**Crosstab**

| | | | GTOTAL1 | | Total |
|---|---|---|---|---|---|
| | | | 1.00 | 2.00 | |
| เลี้ยงดู | แบบเข้มงวด | Count | 18 | 25 | 43 |
| | | % within เลี้ยงดู | 41.9% | 58.1% | 100.0% |
| | | % within GTOTAL1 | 21.7% | 21.6% | 21.6% |
| | แบบปล่อยปละละเลย | Count | 2 | 6 | 8 |
| | | % within เลี้ยงดู | 25.0% | 75.0% | 100.0% |
| | | % within GTOTAL1 | 2.4% | 5.2% | 4.0% |
| | แบบอิสระเสรี | Count | 63 | 85 | 148 |
| | | % within เลี้ยงดู | 42.6% | 57.4% | 100.0% |
| | | % within GTOTAL1 | 75.9% | 73.3% | 74.4% |
| Total | | Count | 83 | 116 | 199 |
| | | % within เลี้ยงดู | 41.7% | 58.3% | 100.0% |
| | | % within GTOTAL1 | 100.0% | 100.0% | 100.0% |

**Chi-Square Tests**

| | Value | df | Asymp. Sig. (2-sided) |
|---|---|---|---|
| Pearson Chi-Square | .964(a) | 2 | .618 |
| Likelihood Ratio | 1.022 | 2 | .600 |
| Linear-by-Linear Association | .044 | 1 | .834 |
| N of Valid Cases | 199 | | |

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 3.34.

**Symmetric Measures**

| | | Value | Approx. Sig. |
|---|---|---|---|
| Nominal by Nominal | Phi | .070 | .618 |
| | Cramer's V | .070 | .618 |
| | Contingency Coefficient | .069 | .618 |
| N of Valid Cases | | 199 | |

a Not assuming the null hypothesis.
b Using the asymptotic standard error assuming the null hypothesis.